arXiv:1302.6193v1 [math.AP] 25 Feb 2013

THE BROKEN RAY TRANSFORM ON THE SQUARE

MARK HUBENTHAL

Abstract. We study a particular broken ray transform on the Euclidean unit square and establish injectivity and stability for C_0^2 perturbations of the constant unit weight. Given an open subset E of the boundary, we measure the attenuation of all broken rays starting and ending at E with the standard optical reflection rule. Using the analytic microlocal approach of Frigyik, Stefanov, and Uhlmann for the X-ray transform on generic families of curves, we show injectivity via a path unfolding argument under suitable conditions on the available broken rays. Then we show that with a suitable decomposition of the measurement operator via smooth cutoff functions, the associated normal operator is a classical pseudo differential operator of order -1 plus a smoothing term with C_0^∞ Schwartz kernel, which leads to the desired result.

1. Introduction

In this article, we are interested in the questions of injectivity and stability for the Euclidean broken ray transform on the unit square in 2-dimensions. Much work has already been done with regard to the attenuated ray transform ([Bal04, Nov02a, Nov02b, BS04, Fin86, Fin04, Nat01]), and for the more general geodesic X-ray transform on smooth manifolds ([FSU08, SU04]), even in the case of generic families of curves. In particular, the main condition for injectivity is, roughly speaking, that the union of the conormal bundles over all available curves must span the cotangent space of the domain (see [FSU08]). Such aforementioned results typically make the assumption that all rays have no conjugate points. In the Riemannian case, this is true if one assumes, for example, that the underlying manifold is simple. But of course such a condition trivially holds in the Euclidean case. However, more recently Stefanov and Uhlmann considered the case of microlocal invertibility for the geodesic ray transform when there are possible isolated conjugate points along certain geodesics, which is the setting for fold caustics ([SU12]).

The main tool used to prove injectivity in [FSU08] for the generalized X-ray transform, and more generally to recover the singularities of an unknown interior distribution f, is analytic microlocal analysis. We employ such ideas heavily in this work. However, it should be stated that other results pertaining to the attenuated ray transform or geodesic ray transform have utilized different methods.

The framework of this paper is of a simpler geometric nature, since few results have been attained for the problem considered so far. Roughly speaking, if (Ω, g) is a compact Riemannian manifold with appropriate conditions, and $E \subset \partial\Omega$ is open, the question is whether the integrals of an unknown function f along all broken rays starting and ending on E uniquely determine f. Unfortunately, reflection tomography of this nature for non-Euclidean metrics has not yet been treated by any previous work. Indeed, the technique of [FSU08] relies on constructing local

---

Date: February 25, 2013.

coordinates about some fixed geodesic γ_0 in order to show the normal operator has nice microlocal properties. But, if one were to reflect from the boundary the family of geodesics associated with such coordinates, one no longer has local coordinates around the reflected segment of γ_0. As such, we will focus just on the Euclidean case, and in particular we will consider the unit square in $\mathbb{R}^2$. It is the author's belief, however, that the results herein are generalizable to at least polyhedral domains which tile $\mathbb{R}^n$, though the computational details may be messy.

There is, however, some work that is directly related to our problem of interest. Eskin proved a uniqueness result for C^∞ unknown functions in 2-dimensions given knowledge of broken rays that reflect off of known convex obstacles in the domain's interior ([Esk04]). There the weight function is taken to be a smooth electric potential $V(x)$. One of the main tools used in the proof is an identity of Mukhometov. Furthermore, the recent work of Ilmavirta in [Ilm12] proves an injectivity result on the disk for open subsets E of the boundary when the unknown function f is uniformly quasianalytic in the angular variable (when written in polar coordinates).

Finally, we mention some recent work of Salo and Kenig, which provides explicit motivation for the consideration of such a broken ray transform in the first place ([KS12]). There the authors consider the recovery of a Schrödinger potential on a particular class of non-compact manifolds from partial Cauchy data. This of course directly relates to the Calderón problem with partial data. It turns out that the question of unique determination of the potential from such partial data would be resolved if one can show injectivity of the broken ray transform considered here, albeit in a more general context.

The structure of this paper goes as follows. In §2 we describe the problem and notations and then state the main results. §3 applies the microlocal ideas of [FSU08] together with the path unfolding technique to show injectivity in the case of 0 attenuation. In §4 we pursue stability of the inverse problem by analyzing the normal operator. To this end, §4.1, 4.2 and 4.3 all deal with computation of the localized normal operator for different choices of neighborhoods of broken rays. Finally, we extend the stability estimate and injectivity to C^2 perturbations of the attenuation σ in §5.

Acknowledgements. This research was conducted with partial support from the Academy of Finland while at the University of Jyväskylä and from the Royal Swedish Academy of Science while at the Mittag-Leffler Institute. The author would like to thank Mikko Salo for introducing him to this problem and for providing helpful comments and feedback.

2. Statement of Main Results

We will state the problem setup and definition in the case of a general domain Ω with smooth boundary. However, the main theorems themselves assume further that Ω is the square. Let Ω be a smooth bounded domain in $\mathbb{R}^n$ and define $\Gamma_\pm = \{(x,\theta) \in \partial\Omega \times \mathbb{S}^{n-1} \mid \pm\nu(x)\cdot\theta > 0\}$ as the set of outgoing and ingoing unit vectors on $\partial\Omega$, respectively. Here $\nu(x)$ is the outward unit normal vector to $\partial\Omega$. We also define, for a general open subset $E \subset \partial\Omega$, the set

$$\Gamma_\pm(E) := \{(x,\theta) \in S\Omega \mid x \in E, \pm\theta\cdot\nu(x) > 0\}. \tag{1}$$

Assume that $\sigma \in C^\infty(\Omega \times \mathbb{S}^{n-1})$ and vanishes near $\partial\Omega$.

We will call any unit speed curve $\gamma_{x,\theta}$ a *regular broken ray* in Ω if

(a) its associated tangent vector field $(\gamma_{x,\theta}(t), \dot{\gamma}_{x,\theta}(t))$ intersects $\Gamma_-(E)$

(b) it consists of finitely many straight line segments $\gamma_{x,\theta,1}, \gamma_{x,\theta,2}, \dots, \gamma_{x,\theta,N}$ before hitting E again

(c) it does not intersect ∂E, and

(d) it obeys the geometrical optics reflection law

$$\dot{\gamma}_{x,\theta,j+1}(0) = -\dot{\gamma}_{x,\theta,j}(L_j) + 2\left(\dot{\gamma}_{x,\theta,j}(L_j) - \nu(\gamma_{x,\theta,j}(L_j)) \cdot \dot{\gamma}_{x,\theta,j}(L_j)\right). \tag{2}$$

Here L_j denotes the length of $\gamma_{x,\theta,j}$ and the subscripts refer to the fact that $(\gamma_{x,\theta}(0), \dot{\gamma}_{x,\theta}(0)) = (x, \theta)$.

We use the notation $T : \Gamma_- \to \Gamma_-$ to denote the billiard map mapping a vector $(x, \theta) \in \Gamma_-$ to $(x', \theta') \in \Gamma_-$, where x' is the intersection point of the geodesic $\gamma_{x,\theta}$ with $\partial\Omega$, and θ' is the reflected direction. We use the maps π_1, π_2 to denote the projections $\Gamma_- \to \partial\Omega$ and $\Gamma_- \to \mathbb{S}^{n-1}$, respectively. The broken ray transform with respect to the accessible set E, denoted by $I_{\sigma,E} : L^2(\Omega) \to L^2(\Gamma_-, d\Sigma)$, is then defined by

$$\begin{aligned}
&I_{\sigma,E} f(\gamma_{x,\theta}) = I_{\sigma,E} f(x,\theta) \\
&:= \sum_{j=0}^{N(x,\theta)} \int_{\mathbb{R}^+} \exp\left(- \sum_{m=0}^{j-1} \int_{\mathbb{R}^+} \sigma(\pi_1 \circ T^m(x,\theta) + \tau \pi_2 \circ T^m(x,\theta), \pi_2 \circ T^m(x,\theta)) \, d\tau \right) \\
&\qquad \cdot \exp\left(- \int_{\mathbb{R}^+} \sigma(\pi_1 \circ T^j(x,\theta) + (t-\tau)\pi_2 \circ T^j(x,\theta), \pi_2 \circ T^j(x,\theta)) \, d\tau \right) \\
&\qquad \cdot f(\pi_1 \circ T^j(x,\theta) + t\pi_2 \circ T^j(x,\theta)) \, dt. \\
&= \sum_{j=0}^{N(x,\theta)} \int_{\mathbb{R}^+} [w_j f] \left(\pi_1 \circ T^j(x,\theta) + t\pi_2 \circ T^j(x,\theta), \pi_2 \circ T^j(x,\theta) \right) dt. \\
&= \sum_{j=0}^{N(x,\theta)} \int_{\mathbb{R}^+} [w_j f] \left(z_j(x,\theta) + t\theta_j(x,\theta), \theta_j(x,\theta) \right) dt.
\end{aligned} \tag{3}$$

for all regular broken rays $\gamma_{x,\theta}$. Here

$$(z_j(x,\theta), \theta_j(x,\theta)) = T^j(x + \tau_-(x,\theta)\theta, \theta), \quad (x,\theta) \in S\Omega \tag{4}$$

where $S\Omega$ is the unit tangent bundle. Also $\tau_\pm(x,\theta) = \min\{t \geq 0 \,|\, x + t\theta \in \partial\Omega\}$ are the positive and negative travel times from a given unit tangent vector $(x,\theta) \in S\Omega$, and $N(x,\theta)$ denotes the number of reflections of the given trajectory before intersecting E. The weight functions w_j on $S\Omega$ are given by

$$\begin{aligned}
w_j(y,\eta) &= \exp\left(- \sum_{m=0}^{j-1} \int_{\mathbb{R}_+} \sigma(z_{m-j} + \tau\theta_{m-j}, \theta_{m-j}) \, d\tau \right) \exp\left(- \int_{\mathbb{R}_+} \sigma(y - \tau\eta, \eta) \, d\tau \right) \\
&= \exp\left(- \sum_{m=1}^{j} \int_{\mathbb{R}_+} \sigma(z_{-m} + \tau\theta_{-m}, \theta_{-m}) \, d\tau \right) \exp\left(- \int_{\mathbb{R}_+} \sigma(y - \tau\eta, \eta) \, d\tau \right)
\end{aligned}$$

The measure $d\Sigma = |\nu(x) \cdot \theta| \, dS(x) \, d\theta$ is the standard measure on Γ_-. It is important to note that because of our assumption that σ vanishes near $\partial\Omega$, this ensures that the weight functions w_j are smooth. The reason is that we can circumscribe Ω with the ball $B = B\left(\left(\frac{1}{2}, \frac{1}{2}\right), \frac{\sqrt{2}}{2}\right)$ and then replace $z_{m-j}(y,\eta) = \pi_1 \circ T^{m-j}(y + \tau_-(y,\eta)\eta, \eta)$ with the last point of ∂B that intersects the line $z_{m-j}(y,\eta) + t\theta_{m-j}(y,\eta)$ for $t \leq 0$. This point is smoothly dependent on $(y,\eta) \in \Omega \times \mathbb{S}^{n-1}$.

Remark 1. Although general polyhedral domains lack obvious reflection rules at the corners (some authors ignore such trajectories and some define them in terms of limits of regular trajectories), the special symmetrical nature of the square does give an unambiguous rule for such reflections. But one still runs into the problem that the composite functions $z_j(x,\theta), \theta_j(x,\theta)$ are not real analytic for (x,θ) associated with a corner point. One simple, albeit more restrictive route, is to assume that the subset E also includes a small neighborhood of each corner. In that case, all broken rays with corner points are just ballistic X-rays. More generally, one can cut out the rays with corner points from the measurement operator. However, if one assumes the attenuation coefficient to vanish in a neighborhood of the boundary, it is still possible to treat rays with corners as we will discuss later.

It is easy to see that $N(x,\theta)$ is piecewise constant, which leads to the unfortunate fact that $I_{\sigma,E}$ in general does not preserve the smoothness of a function. Since we are considering a convex domain (flat) in this work, the discontinuities of $N(x,\theta)$ occur only for $(x,\theta) \in \Gamma_-(E)$ such that $\gamma_{x,\theta}$ has a reflection occurring at some point in ∂E. This is the reason they are not classified as *regular* broken rays. But fortunately, this problem can be averted by omitting such rays from the data with a smooth cutoff.

To eliminate the discontinuities introduced by $N(x,\theta)$ we will use a collection of smooth cutoff functions α_k on $\Gamma_-(E)$ so as to remove the bad parts of $I_{\sigma,E}$. For example, for each point $(x_0,\theta_0) \in \Gamma_-(E)$ where $N(x,\theta)$ is continuous (i.e. γ_{x_0,θ_0} and nearby rays do not touch corner points), we may multiply $I_{\sigma,E}$ by a smooth function $\alpha(x,\theta) \equiv 1$ near (x_0,θ_0) and such that $N(x,\theta)|_{\mathrm{supp}(\alpha)}$ is constant. This localized operator will have a constant number of reflections and thus behaves very much like the usual generalized ray transform, which makes it possible to apply the microlocal analytic techniques of [FSU08]. Of course, additional care is needed to deal with broken rays that contain corner points, which we discuss later. The central idea is that for fixed $(x,\xi) \in T^*\Omega \setminus 0$, any broken ray γ passing through x with direction normal to ξ will allow for the detection of (x,ξ) in $\mathrm{WF}_A(f)$. From hereon we use the notation $\mathrm{WF}_A(f)$ to denote the analytic wavefront set of f.

The main idea we use in order to apply microlocal analytic techniques to the operator is a well-known method of billiards called path unfolding. Essentially, we can unfold any broken ray in Ω into a straight line on a different domain $\widetilde{\Omega}$ defined by reflecting Ω across each of its edges repeatedly to obtain a tiling of $\mathbb{R}^2$. Similarly, we can extend f to a function $\widetilde{f}$ on $\widetilde{\Omega}$ by reflection across $\partial\Omega$. Finally, if $K \subset \Omega$, then we construct $\widetilde{K} \subset \widetilde{\Omega}$ by concatenating reflected copies of K. We will use indices $(l_1,l_2) \in \mathbb{Z}^2$ to denote how many horizontal and vertical reflections a given broken ray undergoes on the unfolded domain $\widetilde{\Omega}$ before returning to the set E. We call this the *reflection signature* of a path. For example, $(-1,2)$ indicates 1 reflection to the left and 2 reflections upwards. The following maps and their respective inverses will be useful when translating between vectors on $T\Omega$ and $T\widetilde{\Omega}$. Given a reflection signature (l_1,l_2), we can compute the reflected version of a point



FIGURE 1. 2-reflected unfolded paths from x to y.

$w \in \Omega$ by

$$\widetilde{w} = R_{l_1,l_2}(w) = \left(l_1 + \frac{1-(-1)^{l_1}}{2} + (-1)^{l_1} w_1, l_2 + \frac{1-(-1)^{l_2}}{2} + (-1)^{l_2} w_2 \right) \tag{5}$$

$$w = R^{-1}_{l_1,l_2}(\widetilde{w}) = \left((-1)^{l_1+1} l_1 + \frac{1-(-1)^{l_1}}{2} + (-1)^{l_1} \widetilde{w}_1,\ (-1)^{l_2+1} l_2 + \frac{1-(-1)^{l_2}}{2} + (-1)^{l_2} \widetilde{w}_2 \right). \tag{6}$$

Similarly, given an angle θ, $S_{l_1,l_2}(\theta)$ is the angle after an (l_1, l_2) reflection.

$$\eta = S_{l_1,l_2}(\theta) = (-1)^{l_1+l_2}\theta + \frac{\pi}{2}\left[1 - (-1)^{l_1}\right] \tag{7}$$

$$\theta = S^{-1}_{l_1,l_2}(\eta) = (-1)^{l_2}\left((-1)^{l_1}\eta + \frac{1-(-1)^{l_1}}{2}\pi \right). \tag{8}$$

Such formulas can be verified with straightforward geometry.

However, applying such unfolding requires significant symmetry conditions on the attenuation σ. First we define an extension of σ on $\widetilde{\Omega} \times \mathbb{S}^1$ by

$$\widetilde{\sigma}(x,\theta) = \sigma(R^{-1}_{l_1,l_2}(x), S^{-1}_{l_1,l_2}(\theta)) \text{ for } x \in [l_1, l_1+1] \times [l_2, l_2+1], \quad (l_1, l_2) \in \mathbb{Z}^2. \tag{9}$$

We then require that $\widetilde{w}(x,\theta) := \exp\left(-\int_0^{L(\gamma_{x,-\theta})} \widetilde{\sigma}(x - \tau\theta, \theta)\, d\tau\right)$ is real analytic for all $(x,\theta) \in \widetilde{\Omega} \times \mathbb{S}^1$ such that the unfolded broken ray through (x,θ) is regular. We've used the notation $L(\gamma_{x,-\theta})$ to mean the length of the half broken ray with initial condition $(x, -\theta)$ before hitting some copy of E in $\widetilde{\Omega}$. Obviously, $\sigma \equiv 0$

works. But such a condition implies that $\widetilde{\sigma}$ has symmetries somewhat like to that of a doubly periodic real analytic function, which would necessarily be constant (the Weierstrass p-function is doubly periodic but has poles of course). Having $\sigma = 0$ near $\partial\Omega$ also allows for $\widetilde{w}$ to be real analytic since then the singularities of $\tau(x, \theta)$ near the corners are masked. Thus, for the purposes of this paper, we will start with the case that the weight function is identically 1 (equivalently $\sigma = 0$), and then prove stability for C^2 perturbations of σ that remain 0 in a neighborhood of $\partial\Omega$.

Note that one can also write $I_{\sigma,E}f$ in the form

$$I_{\sigma,E}f(x, \theta) = \int_0^{L(\gamma_{x,\theta})} \widetilde{w}(x + t\theta, \theta)\widetilde{f}(x + t\theta)\, dt \tag{10}$$

though we will not use it extensively in the subsequent discussion.

For a given $N_{max} \in \mathbb{N}$, we define the *visible set* by

$$\mathcal{M} := \{x \in \Omega \,|\, \forall \xi \in \mathbb{S}^1, (x, \xi) \in N^*\gamma \text{ for some regular broken ray } \gamma \text{ with } N(\gamma) \le N_{max}\}, \tag{11}$$

We can now state the following injectivity result for the 2-dimensional square with Euclidean broken rays:

Theorem 1. *Let $E \subset \partial\Omega$ be open and take $\sigma = 0$. Let $N_{max} \in \mathbb{N}$, let $\mathcal{M}$ be the corresponding visible set, and let $K \Subset \mathcal{M}$. Then $I_{0,E}$ is injective on $L^2(K)$.*

One can also obtain a stability result analogous to that of [FSU08]. To do so, we must first parametrize a family of regular broken rays, having an upper bound on the number of reflections, and whose conormal bundles cover $T^*\Omega$. Fix a positive $N_{max} \in \mathbb{N}$. For each regular broken ray γ_{x_0,θ_0} with $N(x_0, \theta_0) \le N_{max}$, let (l_1, l_2) be it's reflection signature on $\widetilde{\Omega}$. If γ_{x_0,θ_0} has no corner points, then find a maximal connected neighborhood $U \subset \Gamma_-(E)$ containing (x_0, θ_0) such that all broken rays with initial data in U are regular, have no corner points, and have the same reflection signature (l_1, l_2) (See Figure 2). In particular, U itself contains no corner points. We then choose a smooth cutoff function α_k on Γ_- that is equal to 1 on a compact subset of U and vanishes outside of U.

If x_0 is a corner point, then we consider all directed lines $\mathcal{L} = \{(x+t\theta, \theta) \,|\, t \in \mathbb{R}\}$ for (x, θ) in some neighborhood of (x_0, θ_0) such that $N(x, \theta)$ is constant. Then take the intersection of $\mathcal{L}$ with the ball $B = \left(\left(\frac{1}{2}, \frac{1}{2}\right), \frac{\sqrt{2}}{2}\right)$ of radius $\frac{\sqrt{2}}{2}$ whose boundary ∂B circumscribes Ω. We then define a smooth function α on $\partial B \times \mathbb{S}^1$ supported in $\partial B \times \mathbb{S}^1 \cap \mathcal{L}$. We may then extend α to $\mathbb{R}^2 \times \mathbb{S}^1$ to be constant on all directed lines $(x + t\theta, \theta)$. Its restriction to $\Gamma_-(\partial\Omega)$ is the cutoff we seek. We will treat the case that γ_{x_0,θ_0} contains an additional corner point in §4.3.

Let $\alpha = \sum_{k=1}^{N_\alpha} \alpha_k$ be a sum of smooth cutoff functions on $\Gamma_-(E)$ in the sense of the previous paragraph, and consider the regularized operator

$$I_{\sigma,E,\alpha} := \sum_{k=1}^{N_\alpha} \alpha_k I_{\sigma,E}. \tag{12}$$

Then

$$N_{\sigma,E,\alpha} := I^*_{\sigma,E,\alpha} I_{\sigma,E,\alpha} = \sum_{k_1=1,k_2=1}^{K} I^*_{\sigma,E}\alpha_{k_1}\alpha_{k_2} I_{\sigma,E}. \tag{13}$$



FIGURE 2. A typical beam of unfolded broken rays which follow the same reflection sequence before hitting E. The beam terminates in the part of $\widetilde{\Omega}$ with reflection signature $(h_4, v_4) = (2, 2)$.

The transpose is taken with respect to the standard measure $|\nu(x) \cdot \theta| \, dS(x) \, d\theta$ on $\Gamma_-(E)$. Note that we may as well assume that the α_k functions have disjoint support, since it won't change the kind of terms present in the normal operator (if two such cutoff functions α_{k_1} and α_{k_2} have overlapping support, then their product is also smooth with the supported broken rays localized just as in the previous construction).

We also define the *microlocally visible set* by

$$\mathcal{M}' := \{(z,\xi) \in T^*\Omega \,|\, (z,\xi) \in N^*\gamma_{x,\theta} \text{ for some } (x,\theta) \text{ with } \alpha(x,\theta) > 0\}. \tag{14}$$

Remark 2. One reason we need to impose an upper bound on the number of reflections is that as N increases, the open level sets of $N(X,\theta)$ shrink to zero. Similarly, the neighborhoods in Γ_- on which we have constant reflection signature (l_1, l_2) become infinitesimal as $|l_1| + |l_2| \to \infty$. Thus α will not be smooth unless we apply some threshold.

We have the following stability result analogous to that in [FSU08].

Theorem 2. (a) *Fix σ that is 0 in a neighborhood of $\partial\Omega$, and choose $\alpha \in C^\infty(\Gamma_-)$ as above with reflection threshold N_{max}. Fix a set $K \Subset \mathcal{M}$ compactly contained in the visible set $\mathcal{M}$. If $I_{\sigma,E,\alpha}$ is injective on $L^2(K)$, then*

$$\frac{1}{C}\|f\|_{L^2(K)} \le \|N_{\sigma,E,\alpha} f\|_{H^1(\Omega)} \le C\|f\|_{L^2(K)}. \tag{15}$$

(b) *Let α^0 be as above related to some fixed σ_0. Assume that I_{σ_0,E,α^0} is injective on $L^2(K)$. Then estimate (15) remains true for (σ,α) in a small C^2 neighborhood of (σ_0,α^0) such that $\sigma \equiv 0$ near $\partial\Omega$, and with a uniform constant $C > 0$.*

3. INJECTIVITY OF $I_{0,E}$ FOR THE SQUARE

For the 2-dimensional square, we will be able to establish injectivity of the broken ray transform for the unknown function f supported on some subset of the domain that depends on E. We will be able to detect covectors from $N^*\gamma_{x,\theta}$ in $\mathrm{WF}_A(f)$ for (most) $(x,\theta) \in \Gamma_-(E)$. The idea for the proof of the theorem is heavily borrowed from the microanalytic techniques utilized in [FSU08]. We will need to assume that f is compactly supported within Ω, which in particular will allow us to treat the reflected ray as a smooth regular curve in the sense of [FSU08] since the integrand vanishes near the points of reflection. By using the path unfolding approach, we can relatively easily obtain the following microlocal result.

Proposition 1. *Let $f \in \mathcal{D}'(K)$ for $K \Subset \Omega$. Suppose $I_{0,E}f(x,\theta) = 0$ for all (x,θ) in a small neighborhood of $(x_0,\theta_0) \in \Gamma_-(E)$ where γ_{x_0,θ_0} is a regular broken ray. Then $\mathrm{WF}_A(f) \cap N^*\gamma_{x_0,\theta_0} = \emptyset$.*

Proof. We need to construct coordinates near the broken ray γ_0. To do this, we let $\{x_{0,1},\dots,x_{0,N}\}$ be the ordered reflection points on $\partial\Omega$ for γ_0, where $N = N(x_0,\theta_0)$. We then construct a new domain $\widetilde{\Omega}$ consisting of N reflected copies of Ω and glued together with the original Ω along the edges that contain each $x_{0,j}$, $1 \le j \le N$. Specifically, let P_m for $1 \le m \le 4$ be the lines in $\mathbb{R}^2$ which bound the sides of Ω in some order. Let $\{m_j\}_{j=1}^N$ be the indices corresponding to which edge of $\partial\Omega$ that $x_{0,j}$ belongs. We iteratively construct $\widetilde{\Omega}$ by starting with $x_{0,1}$ and reflecting Ω across edge P_{m_1}. Then glue the two copies of Ω together along P_{m_1}. Continuing in this way, we obtain a space consisting of $N+1$ copies of Ω, each reflected along a specific direction. If γ_0 contains a corner point, we can add additional reflected copies of Ω as necessary to accommodate nearby broken rays. We also obtain a distribution $\widetilde{f} \in \mathcal{D}'(\widetilde{\Omega})$ and an unfolded version of γ_0, denoted by $\widetilde{\gamma_0}$, using the same process. Note that by reversing the reflection sequence, any covector $(\widetilde{z},\widetilde{\xi}) \in T^*\widetilde{\Omega}$ corresponds to a unique covector $(z,\xi) \in T^*\Omega$.

Since the edges of Ω are flat, it follows that $\widetilde{\gamma_0}$ is a straight line. The openness of E and the assumptions on (x_0,θ_0) ensure that for (x,θ) in a neighborhood $U \subset \Gamma_-(E)$ of (x_0,θ_0), the unfolded paths $\widetilde{\gamma}_{x,\theta}$ lie in the same extended space $\widetilde{\Omega}$ and avoid ∂E. Now choose a point p_0 on the line $x_0 + t\theta_0$ for $t < 0$ so that $p_0 \notin \Omega$. Define $x = p_0 + t\theta$. Then (θ,t) are local coordinates near any point of $\widetilde{\Omega} \cap \widetilde{\gamma}_{x_0,\theta_0}$. We can assume without loss of generality that θ_0 corresponds to the vector $(0,1)$. Writing $x = (x^1,x^2) = (\theta^1,t)$, we have the coordinates

$$U = \{x \,|\, |x^1| < \epsilon,\, l^- < t < l^+\} \subset \Omega.$$

Since f is compactly supported inside of Ω, we may shrink ϵ as necessary so that we can take l^-, l^+ to be constant. Now as in the proof of Proposition 1 in [FSU08], we let $(z_0,\xi_0) \in N^*\gamma_0$. Then consider the corresponding reflected covector via path unfolding given by $(\widetilde{z}^0,\widetilde{\xi_0}) \in N^*\widetilde{\gamma_0}$. The argument of [FSU08] shows that $(\widetilde{z}^0,\widetilde{\xi_0}) \notin \mathrm{WF}_A(\widetilde{f})$. By undoing the reflection process again, we have that $(z_0,\xi_0) \notin \mathrm{WF}_A(f)$. $\square$

Proof of Theorem 1. Suppose $I_{\sigma,E}f = 0$. By Theorem 1 we have that f is analytic on K. Since $\mathrm{supp}(f) \subset K$, we have that f can be extended to an entire function and hence must be identically zero. $\square$

Remark 3. In two dimensions, Theorem 1 is not so remarkable since in this case every unit vector has a unique unit normal vector up to sign. For example, in the case of the square Theorem 1 gives injectivity on the whole domain if E contains two adjacent (closed) edges. However, one must be careful since it fails for E consisting of two opposite edges.

4. Stability

In order to prove the stability result for $I_{\sigma,E}$ with σ a C^2 perturbation of 0 that vanishes near $\partial\Omega$, we will first compute its normal operator. Suppose there is an upper limit N_{max} on the number of reflections we want to allow in the data, which is implicit in the choice of cutoff functions $\alpha_k(x,\theta)$. We construct the modified operator

$$I_{\sigma,E,\alpha}f(x,\theta) := \sum_{k=1}^{N_\alpha} \alpha_k(x,\theta) I_{\sigma,E}f(x,\theta), \quad (x,\theta) \in \Gamma_-(E). \tag{16}$$

We will always suppose that $K \Subset \mathcal{M}$ is fixed, where $\mathcal{M} \subset \Omega$ is the visible set. We will prove the following proposition regarding the structure of $N_{\sigma,E,\alpha}$:

Proposition 2. *$N_{\sigma,E,\alpha} = N_{\sigma,E,\alpha,ballistic} + N_{\sigma,E,\alpha,reflect}$ where $N_{\sigma,E,\alpha,ballistic}$ is a classical pseudo differential operator of order -1, elliptic on $\mathcal{M}$, and $N_{\sigma,E,\alpha,reflect}$ is an operator with $C_0^\infty(\Omega\times\Omega)$ Schwartz kernel. Thus there exists a classical pseudo differential operator Q in Ω of order 1 such that*

$$QN_{\sigma,E,\alpha}f = f + QN_{\sigma,E,\alpha,reflect}f + S_1 \tag{17}$$

for any $f \in \mathcal{D}'(K)$, and an operator S_1 with $C_0^\infty(\Omega \times \Omega)$ Schwartz kernel.

4.1. Computing the Normal Operator .

Recall that given $j \in \mathbb{Z}$ we have defined the variables z_j, θ_j depending on $x \in \Omega$, $\theta \in \mathbb{S}^{n-1}$ by

$$(z_j, \theta_j) := T^j(x + \tau_-(x,\theta)\theta,\ \theta) \tag{18}$$

Notice that $(z_0,\theta_0) = (x + \tau_-(x,\theta)\theta, \theta)$.

Choose a cutoff function $\alpha(x,\theta)$ on Γ_- such that $\alpha I_{\sigma,E}$ has a constant number of reflections $N \geq 0$ on its support. Later we will consider broken rays with corner points, but then the number of reflections is not constant, so it must be treated carefully. The adjoint to $\alpha I_{\sigma,E}$ can be computed as follows

$$\begin{aligned} &\int_{\Gamma_-} \alpha(x,\theta)[I_{\sigma,E}f](x,\theta)g(x,\theta)\, d\Sigma \\ &\quad = \int_{\Gamma_-} \alpha(x,\theta)g(x,\theta) \sum_{j=0}^{N} \int_{\mathbb{R}^+} w_j(z_j + t\theta_j, \theta_j) f(z_j + t\theta_j) |\nu(x)\cdot\theta|\, dt\, dS(x)\, d\theta \end{aligned} \tag{19}$$

For each j we make the change of variables $(x,\theta,t) \mapsto (y,\eta)$ where $(y,\eta) := (z_j(x,\theta) + t\theta_j(x,\theta), \theta_j(x,\theta))$. Thus

$$\begin{aligned} (x,\theta) &= (z_{-j}(y,\eta), \theta_{-j}(y,\eta)) \\ t &= -\tau_-(y,\eta). \end{aligned} \tag{20}$$

Intuitively, in moving from (y,η) to $(z_{-j}(y,\eta),\theta_{-j}(y,\eta))$, we first project from the point y to the boundary along the direction $-\eta$ and then trace backwards along the trajectory for the previous j reflections. We compute

$$\begin{aligned}&\Big(z_m(x,\theta)+s\theta_m(x,\theta),\theta_m(x,\theta)\Big)\\&=\Big(z_m\Big(z_{-j}(y,\eta),\theta_{-j}(y,\eta)\Big)+s\theta_m\Big(z_{-j}(y,\eta),\theta_{-j}(y,\eta)\Big),\theta_m\Big(z_{-j}(y,\eta),\theta_{-j}(y,\eta)\Big)\Big)\\&=(z_{m-j}(y,\eta)+s\theta_{m-j}(y,\eta),\,\theta_{m-j}(y,\eta)).\end{aligned}$$

Given a fixed number of reflections j, it is clear that each point $(y,\eta)\in\Omega\times\mathbb{S}^{n-1}$ corresponds to a unique point $(x,\theta)\in\Gamma_-$. Thus, the previously described change of variables is a diffeomorphism. The resulting integral is over the domain $\Omega\times\mathbb{S}^{n-1}$. We have

$$\sum_{j=0}^{N}\int_{\Omega\times\mathbb{S}^{n-1}}\alpha(z_{-j}(y,\eta),\theta_{-j}(y,\eta))g(z_{-j}(y,\eta),\theta_{-j}(y,\eta))w_j(y,\eta)f(y)J_j(y,\eta)\,dy\,d\eta,$$

where

$$J_j(y,\eta)=\left|\frac{\partial x\partial\theta\partial t}{\partial y\,\partial\eta}\right||\nu(z_{-j}(y,\eta))\cdot\theta_{-j}(y,\eta)|.$$

Since the billiard map T preserves the symplectic form $|\nu(x)\cdot\theta|dx\wedge d\theta$ on Γ_-, it follows that $J_j(y,\eta)=1$. We stress that this is even true for domains with piecewise smooth boundary as is the case for the square, by measure theoretic arguments (see [Tab] §1.7). One can verify that the Jacobian factors above cancel by first making the change of variables $(x,\theta)\to T^j(x,\theta)$ in (19), using the invariance of the symplectic form $|\nu(x)\cdot\theta|dx\wedge d\theta$ under T. Then change to (y,η) coordinates on $\Omega\times\mathbb{S}^{n-1}$ using the same approach as in (Theorem 1 (b), [SU08]). Thus we have

$$(\alpha I_{\sigma,E})^*g(x)=\sum_{j=0}^{N}\int_{\mathbb{S}^{n-1}}[\alpha g](z_{-j}(x,\theta),\theta_{-j}(x,\theta))w_j(x,\theta)\,d\theta\tag{21}$$

We are now ready to compute the normal operator $I^*_{\sigma,E}I_{\sigma,E}$. By definition

$$\begin{aligned}&I^*_{\sigma,E,\alpha}I_{\sigma,E,\alpha}f(x)\\&=\sum_{j_1=0}^{N}\int_{\mathbb{S}^{n-1}}|\alpha(z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))|^2I_{\sigma,E}f(z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))w_{j_1}(x,\theta)\,d\theta\\&=\sum_{j_1=0}^{N}\sum_{j_2=0}^{N}\int_{\mathbb{S}^{n-1}}\int_{\mathbb{R}_+}|\alpha(z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))|^2\,w_{j_1}(x,\theta)\\&\quad\cdot w_{j_2}(z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta),\theta_{j_2-j_1}(x,\theta))\,f\left(z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta)\right)\,dt\,d\theta\\&=\sum_{j_1=0}^{N}\sum_{j_2=0,j_2\neq j_1}^{N}\int_{\mathbb{S}^{n-1}}\int_{\mathbb{R}_+}w_{j_1}(x,\theta)\,w_{j_2}(z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta),\theta_{j_2-j_1}(x,\theta))\\&\quad\cdot|\alpha(z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))|^2\,f\left(z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta)\right)\,dt\,d\theta\\&+\sum_{j_1=0}^{N}\int_{\mathbb{S}^{n-1}}\int_{\mathbb{R}_+}w_{j_1}(x+t\theta,\theta)w_{j_1}(x,\theta)\,|\alpha(z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))|^2\,f\left(x+\tau_-(x,\theta)\theta+t\theta\right)\,dt\,d\theta\end{aligned}$$

(22)

$$=:N_{\sigma,E,\alpha,reflect}f(x)+N_{\sigma,E,\alpha,ballistic}f(x).$$

It follows that the full normal operator $N_{\sigma,E,\alpha}$, corresponding to the partition by smooth cutoff functions α_k, is given by the sum

$$\begin{aligned}
&N_{\sigma,E,\alpha} f(x) \\
&= \sum_{k=1}^{N_\alpha} \sum_{j_1=0}^{N_k} \sum_{j_2=0}^{N_k} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}_+} |\alpha_k(z_{-j_1}, \theta_{-j_1})|^2 w_{j_1}(x,\theta)\, w_{j_2}\left(z_{j_2-j_1} + t\theta_{j_2-j_1}, \theta_{j_2-j_1}\right) \\
&\cdot f\left(z_{j_2-j_1} + t\theta_{j_2-j_1}\right)\, dt\, d\theta
\end{aligned}$$

We pick off the ballistic terms where $j_1 = j_2$ to decompose $N_{\sigma,E,\alpha} f$ as a sum

$$N_{\sigma,E,\alpha} = N_{\sigma,E,\alpha,ballistic} + N_{\sigma,E,\alpha,reflect}$$

with

$$\begin{aligned}
&N_{\sigma,E,\alpha,ballistic} f(x) \\
&= \sum_{k=1}^{N_\alpha} \sum_{j=0}^{N_k} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}_+} |\alpha_k(z_{-j}, \theta_{-j})|^2 w_j(x,\theta)\, w_j\left(x + (\tau_-(x,\theta) + t)\theta, \theta\right) \\
&\qquad \cdot f\left(x + (\tau_-(x,\theta) + t)\theta\right)\, dt\, d\theta \\
&= \sum_{k=1}^{N_\alpha} \sum_{j=0}^{N_k} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}} |\alpha_k(z_{-j}, \theta_{-j})|^2 w_j(x,\theta)\, w_j(x + t\theta, \theta)\, f(x + t\theta)\, dt\, d\theta \\
&= \sum_{k=1}^{N_\alpha} \sum_{j=0}^{N_k} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}+} \left[|\alpha_k(z_{-j}(x,\cdot)\theta_{-j}(x,\cdot))(\theta)|^2 w_j(x,\cdot)\, w_j(x + t\theta, \cdot)\right]_{even}(\theta) \\
&\qquad \cdot f(x + t\theta)\, dt\, d\theta
\end{aligned}$$

and

$$\begin{aligned}
N_{\sigma,E,\alpha,reflect} f(x) = \sum_{k=1}^{N_\alpha} \sum_{j_1=0}^{N_k} \sum_{j_2=0, j_2 \neq j_1}^{N_k} \int_{\mathbb{S}^{n-1}} \int_{\mathbb{R}_+} |\alpha_k(z_{-j_1}, \theta_{-j_1})|^2 \\
\cdot w_{j_1}(x,\theta)\, w_{j_2}\left(z_{j_2-j_1} + t\theta_{j_2-j_1}, \theta_{j_2-j_1}\right) f\left(z_{j_2-j_1} + t\theta_{j_2-j_1}\right)\, dt\, d\theta.
\end{aligned}$$

The notation $[g(x,\cdot)]_{even}(\theta)$ is the even part of g in the variable θ, defined by $[g(x,\cdot)]_{even}(\theta) = \frac{1}{2}[g(x,\theta) + g(x,-\theta)]$.

By Lemma 2 of [FSU08] $N_{\sigma,E,\alpha,ballistic}$ is a classical pseudo differential operator of order -1 with principal symbol

$$a_0(x,\xi) = 2\pi \sum_{k=0}^{K} \sum_{j=0}^{N_k} \int_{\theta \in \mathbb{S}^{n-1},\, \theta\cdot\xi = 0} |\alpha_k(z_{-j}, \theta_{-j})|^2 \left|w_j(x,\theta)\right|^2\, d\theta \tag{23}$$

The bound on the number of reflections N_{max} ensures that the integral kernel has a positive lower bound on the set of all covectors (x,ξ) such that it is non vanishing at some $\theta \in \mathbb{S}^{n-1}$ normal to ξ. Clearly, $N_{\sigma,E,\alpha,ballistic}$ is also elliptic on the set $\mathcal{M}'$.

4.2. **Simplifying** $N_{\sigma,E,\alpha,reflect}$ **for the Square .** For the reflected part of the normal operator, our goal will be to make a change of variables so that $f(z_{j_2-j_1} + t\theta_{j_2-j_1})$ becomes $f(y)$, with the hope that what we obtain is either a weakly singular integral operator (i.e. smoothing of order -1) or an operator with C_0^∞ Schwartz kernel. It turns out that the latter will hold. This technique is very much inspired by the kind of substitution used in [SU08, Hub11], and also in [FSU08] for the non-Euclidean case. For now, we will still be working with the part of the operator that avoids broken rays with corner points.

Let $(x,\theta) \in \Omega \times \mathbb{S}^1$, suppose we have some smooth cutoff α on $\Gamma_-(E)$. By construction, all broken rays originating from the support of α have the same sequence of reflection signatures. Let $\{(h_1,v_1),\ldots,(h_N,v_N)\}$ be the ordered set of all such pairs (see Figure 2). Let $z = z_{j_2-j_1}(x,\theta)$ (i.e. z is the j_2^{th} reflection point on $\partial\Omega$ of the broken ray starting at $(z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta)))$. Then $(h_{j_2}-h_{j_1}, v_{j_2}-v_{j_1})$ is the reflection signature from x to z, and so we consider

$$\widetilde{z} := R_{h_{j_2}-h_{j_1},v_{j_2}-v_{j_1}}(z) = (\widetilde{z}^1,\widetilde{z}^2)$$

as the associated unfolded coordinates of z. Up to a rotation of Ω, there are essentially two possibilities depending on whether $\widetilde{z}$ lies on a top or right edge of a reflected copy of the square. We will use θ to denote an angle value in radians and $\widehat{\theta} = (\cos\theta,\sin\theta)$ to denote its associated direction vector. Furthermore, if $x \in \mathbb{R}^2$, we define $x^\perp = (-x^2,x^1)$. For simplicity of notation, we let $(l_1,l_2) = (h_{j_2}-h_{j_1}, v_{j_2}-v_{j_1})$.

We have one of the following equations for $\widetilde{z}$ as a function of the starting point x with direction θ:

$$\widetilde{z} = (\widetilde{z}^1, l_2) = (x^1 + (l_2 - x^2)\cot\theta, l_2) \tag{24}$$

$$\widetilde{z} = (l_1, \widetilde{z}^2) = (l_1, x^2 + (l_1 - x^1)\tan\theta). \tag{25}$$

In the integral expression of the normal operator, we first make the substitution $\eta = S_{l_1,l_2}(\theta)$, which is the reflected angle after applying the billiard map $j_2 - j_1$ times to $(x + \tau_-(x,\widehat{\theta})\widehat{\theta}, \widehat{\theta})$. Note that $|l_1| + |l_2| = j_2 - j_1$. Obviously, the Jacobian determinant of this transformation is 1. We also rewrite $z_{j_2-j_1}(x,\widehat{\theta}) = \pi_1 \circ T^{j_2-j_1}(x + \tau_-(x,\widehat{\theta})\widehat{\theta}, \widehat{\theta})$ as the function $z(x,\eta)$. In particular, we may have either

$$\begin{aligned} z(x,\eta) &= R^{-1}_{l_1,l_2}\left(x^1 + (l_2 - x^2)\cot\left(S^{-1}_{l_1,l_2}(\eta)\right), \quad l_2\right) \\ &= \left((-1)^{l_1+1}l_1 + \frac{1-(-1)^{l_1}}{2} + (-1)^{l_1}\left[x^1 + (l_2 - x^2)\cot\left(S^{-1}_{l_1,l_2}(\eta)\right)\right], \quad \frac{1-(-1)^{l_2}}{2}\right) \end{aligned} \tag{26}$$

or

$$\begin{aligned} z(x,\eta) &= R^{-1}_{l_1,l_2}\left(l_1, \quad x^2 + (l_1 - x^1)\tan\left(S^{-1}_{l_1,l_2}(\eta)\right)\right) \\ &= \left(\frac{1-(-1)^{l_1}}{2}, \quad (-1)^{l_2+1}l_2 + \frac{1-(-1)^{l_2}}{2} + (-1)^{l_2}\left[x^2 + (l_1 - x^1)\tan\left(S^{-1}_{l_1,l_2}(\eta)\right)\right]\right). \end{aligned} \tag{27}$$

Thus we have an integrand involving $f(z(x,\eta) + t\widehat{\eta})$, where $\widehat{\eta} = (\cos\eta,\sin\eta)$. The next step is to make the substitution $y = z(x,\eta) + t\widehat{\eta}$ (see Figure 3). We compute it's Jacobian determinant to be

$$\left|\frac{\partial y}{\partial\eta\partial t}\right| = |(-1)^{l_2}(l_2 - x^2)\csc\eta + t| = |(l_2 - x^2)\csc\theta + t| = |\widetilde{y} - \widetilde{z}|\left|\frac{l_2 - x^2}{\widetilde{y}^2 - \widetilde{z}^2} + 1\right|$$

$$= |\widetilde{y} - \widetilde{z}|\left|\frac{\widetilde{y}^2 - x^2}{\widetilde{y}^2 - l_2}\right| \tag{28}$$


$\widetilde{y}$
$\widetilde{z}$
$(\widetilde{x}, S_{1,1}^{-1}(\theta))$
y
z
(x,θ)
$(z_{-2}(x,\theta),\theta_{-2}(x,\theta))$

FIGURE 3. The basic setup for how we define coordinates in the integral formula for $N_{\sigma,E,\alpha}f$. Here $j_2 = 4$ and $j_1 = 2$.

if $z(x,\eta)$ is given by (26). Similarly, we obtain

$$\left|\frac{\partial y}{\partial\eta\partial t}\right| = |(-1)^{l_1}(l_1 - x^1)\sec\eta + t| = |\widetilde{y} - \widetilde{z}|\left|\frac{\widetilde{y}^1 - x^1}{\widetilde{y}^1 - l_1}\right|$$

when $z(x,\eta)$ is given by (27). Here we've used the fact that $\sec\eta = (-1)^{l_1}\sec\theta$ and $\csc\eta = (-1)^{l_2}\csc\theta$. In order to fully carry out the substitution in the integral, we need to write z as a function of x and y. We will need the auxiliary equation

$$(\widetilde{y} - \widetilde{z}) \cdot (\widetilde{z} - x)^{\perp} = 0, \tag{29}$$

which is a consequence of the reflection law. In the case where $\widetilde{z}^2$ is undetermined, we have $\widetilde{z}^1 = l_1$ and then

$$\widetilde{z}^2 - \widetilde{y}^2 = \frac{(\widetilde{y}^2 - x^2)(l_1 - \widetilde{y}^1)}{\widetilde{y}^1 - x^1}. \tag{30}$$

Similarly, if $\widetilde{z}^1$ is undetermined then

$$\widetilde{z}^1 - \widetilde{y}^1 = \frac{(\widetilde{y}^1 - x^1)(l_2 - \widetilde{y}^2)}{\widetilde{y}^2 - x^2}. \tag{31}$$

In the given case $\widetilde{z}^1 = l_1$, the Jacobian determinant $\left|\frac{\partial y}{\partial\eta\partial t}\right|$ simplifies to

$$\begin{aligned} |\widetilde{y} - \widetilde{z}|\left|\frac{\widetilde{y}^1 - x^1}{\widetilde{y}^1 - l_1}\right| &= \sqrt{(\widetilde{y}^1 - \widetilde{z}^1)^2 + (\widetilde{y}^2 - x^2)^2\frac{(l_1 - \widetilde{y}^1)^2}{(\widetilde{y}^1 - x^1)^2}}\left|\frac{\widetilde{y}^1 - x^1}{\widetilde{y}^1 - l_1}\right| \\ &= |\widetilde{y} - x|. \end{aligned}$$

The same holds when $\widetilde{z}^2 = l_2$. Conveniently, the $|\widetilde{y} - x|$ is strictly positive for all $y, x \in \mathrm{supp}(f)$ so long as l_1 or l_2 is nonzero (i.e. $j_2 - j_1 \neq 0$). Thus the reciprocal factor $\frac{1}{|\widetilde{y}-x|}$ resulting from the change of variables is smooth and bounded as a function of y and x. Using again the fact that $\cos\eta = (-1)^{l_1}\cos\theta$, $\sin\eta = (-1)^{l_2}\sin\theta$ we have that

$$\widehat{\eta} = ((-1)^{l_1}\cos\theta, (-1)^{l_2}\sin\theta) = \frac{1}{|\widetilde{y} - x|}((-1)^{l_1}(\widetilde{y}^1 - x^1), (-1)^{l_2}(\widetilde{y}^2 - x^2)).$$

Altogether, we obtain that $N_{\sigma,E,\alpha,reflect}f(x)$ consists of a sum of terms of the form

$$\int_{\mathbb{R}^2} |\alpha_k(T^{-j_2}(y + \tau_-(y, \widehat{\eta})\widehat{\eta}, \widehat{\eta}))|^2 w_{j_1}\left(x, \frac{\widetilde{y} - x}{|\widetilde{y} - x|}\right) w_{j_2}(y, \widehat{\eta})\, f(y)\, \frac{dy}{|\widetilde{y} - x|} \tag{32}$$

Evidently, $\frac{1}{|\widetilde{y}-x|}$ is a smooth non-vanishing function of x and y since $|\widetilde{y} - x| \geq d(\mathrm{supp}(f), \partial\Omega) > 0$.

Note that one must be very careful in the above formulation in order to show that the integrand in (32) is smooth where it is supported. Consider the case if α_k is non vanishing for a broken ray which starts at a corner point, but still none of the broken rays in the support of α contain another corner point after some positive number of reflections. Within the factor of (32) given by

$$|\alpha_k(T^{-j_2}(y + \tau_-(y, \widehat{\eta})\widehat{\eta}, \widehat{\eta}))|^2,$$

we see that $T^{-j_2}(y+\tau_-(y,\widehat{\eta})\widehat{\eta}, \widehat{\eta})$ fails to be smooth for (y, η) that are pulled back to this corner point. However, as previously discussed, the construction of α_k ensures that it is smooth on $\mathbb{R}^2 \times \mathbb{S}^1$ and recall that $B = B\left(\left(\frac{1}{2}, \frac{1}{2}\right), \frac{\sqrt{2}}{2}\right)$ is the ball that circumscribes Ω. By a possible rotation of Ω, we may as well assume that h_j, v_j are all nonnegative. Clearly each sequence $\{h_j\}$, $\{v_j\}$ is also nondecreasing in this case. Next define the function $q_j(y, \eta)$ for $1 \leq j \leq N$ as the *final* intersection point of the line $R_{h_j,v_j}(y) + t\widehat{S^{-1}_{h_j,v_j}(\eta)}$ with ∂B for $t \leq 0$. We also define $q_0(y, \eta) = y + \tau_{-,\partial B}(y, \eta)\widehat{\eta}$ where $\tau_{-,\partial B}(y, \widehat{\eta})$ is the negative distance to ∂B from y in the direction $-\widehat{\eta}$, defined for $(y, \widehat{\eta}) \in B \times \mathbb{S}^1$. We can then replace $\alpha_k(T^{-j_2}(y + \tau_-(y, \widehat{\eta})\widehat{\eta}, \widehat{\eta}))$ with

$$\alpha_k(q_{j_2}(y, \widehat{\eta}),\, \widehat{S^{-1}_{h_{j_2},v_{j_2}}(\eta)}) \tag{33}$$

Clearly, (33) depends smoothly on y, η.

We observe that the definitions of w_{j_1}, w_{j_2} also involve τ_-, which is not smooth on $\Omega \times \mathbb{S}^1$. However, since σ is compactly supported inside of Ω on some smooth subdomain, we can modify τ_- inside of w_{j_1}, w_{j_2} as necessary to be smooth without changing the integrals. In particular, we can use the functions q_j to define the weight functions instead of the billiard map T and boundary distance function τ_-. Therefore, (32) is a smoothing integral operator.

4.3. **Incorporating Corners.**

While the corners of $\partial\Omega$ have no well-defined normal vector, we can make sense of a broken ray that touches a corner point by realizing it as a limit of broken rays that avoid the corner. This is described in detail in the 2-dimensional case in the work of Eskin in [Esk04]. Essentially, in 2-dimensions there are two reflection directions at a corner point that can be obtained as a limit of broken rays starting at some fixed point $x \in \Omega$ and with reflection point approaching the given corner. In higher dimensions, one can obtain a cone of possible reflection directions, which is thus a hypersurface. However, for the square (or even the n-cube), these limiting rays all conveniently coincide. To see this, let $x \in \Omega$ and let c be a corner point. Define θ_c as the angle between $c - x$ and the horizontal axis. For θ close to θ_c, the broken ray from x in the direction $\widehat{\theta}$ will intersect twice near c and return at an angle $\pi + \theta$ with the positive horizontal direction (i.e. it will return in the direction $-\widehat{\theta}$). Therefore, the limit of such rays shows that γ_{x,θ_c} unambiguously reflects back towards x. This is also consistent with what one expects from path unfolding.

Lemma 1. *A regular broken ray starting from E contains at most two corner points, and if it has two, then the first one is its starting point.*

Proof. Suppose we have a broken ray γ_{x_0,θ_0} with two distinct corner points. Since reflections from corners return in the opposite direction from the direction of incidence, any broken ray between them will repeat periodically. Thus either one of the corner points lies in E or there is a point along the broken ray between them that lies in E. But this implies the broken ray would have terminated before hitting both corners. $\square$

We will use $c \neq x_0$ to refer to a corner in the broken ray if it has one, and we will allow for the possibility that x_0 itself it a corner point. However, as previously discussed, x_0 being a corner point does not present any problem with regard to the smoothness of the integrands involved in the expansion of the localized normal operator (see the discussion following (32)). As before, let $K \Subset \Omega$ be the set on which we assume f has support.

So given our broken ray γ_{x_0,θ_0}, we assume it passes through a corner point $c \neq x_0$ at the m^{th} reflection, where $1 \leq m \leq N-1$, and $N-1$ is the total number of reflections of the curve. Choose a neighborhood U consisting of $(x,\theta) \in \Gamma_-(E)$ that are sufficiently close to (x_0,θ_0) in the following sense. At the m^{th} reflection, most nearby broken rays to γ_{x_0,θ_0} will undergo a reflection along a very short segment near the corner on which f vanishes and $\sigma = 0$. In particular, such segments do not affect the weights w_j and so we will be able to discard them. Thus, we choose U small enough so that at the mth reflection, all nearby broken rays to γ_{x_0,θ_0} either pass through the corner c or hit close enough to the corner so as to reflect into a short path that contributes nothing to the integral. Note that for such broken rays from U passing very near to the corner c (but not actually touching it), after the $(m+1)^{\text{th}}$ reflection they will be oriented in a direction opposite their respective directions during the $(m-1)^{\text{th}}$ segment. We also choose U small enough so as not to introduce any new corner points among the included broken rays.

Given such an open set U, we let $\mathbb{1}_U(x,\theta)$ be a function on $\mathbb{R}^2 \times \mathbb{S}^1$ which is constant on lines and satisfies

(34)
$$\mathbb{1}_U(x,\theta) = \begin{cases} 1 & \text{if } (\gamma_{x,\theta}(t), \dot{\gamma}_{x,\theta}(t) \text{ passes through } U \text{ and intersects the corner } c, \\ 0 & \text{otherwise} \end{cases}$$

Also let $\alpha(x,\theta)$ be a smooth cutoff function in $\mathbb{R}^2\times\mathbb{S}^1$ which vanishes outside the set of lines associated with U. We write the localized broken ray transform as

$$\begin{aligned} I_{\sigma,E,\alpha}f(x,\theta) &= \sum_{j=0}^{N}\int_{\mathbb{R}^+}(1-\mathbb{1}_U(x,\theta))\alpha(x,\theta)[w_jf](z_j(x,\theta)+t\theta_j(x,\theta),\theta_j(x,\theta))\,dt \\ &\quad + \sum_{j=0}^{N-1}\int_{\mathbb{R}^+}\mathbb{1}_U(x,\theta)\alpha(x,\theta)[w_jf](z_j(x,\theta)+t\theta_j(x,\theta),\theta_j(x,\theta))\,dt. \end{aligned}\tag{35}$$

We then have

$$\begin{aligned} I^*_{\sigma,E,\alpha}g(x) &= \sum_{j=0}^{N}\int_{\mathbb{S}^1}[(1-\mathbb{1}_U)\alpha g](z_{-j}(x,\theta),\theta_{-j}(\theta))w_j(x,\theta)\,d\theta \\ &\quad + \sum_{j=0}^{N-1}\int_{\mathbb{S}^1}[\mathbb{1}_U\alpha g](z_-(x,\theta),\theta_{-j}(x,\theta))w_j(x,\theta)\,d\theta. \end{aligned}\tag{36}$$

Finally, the normal operator is given by

$$\begin{aligned} I^*_{\sigma,E,\alpha}I_{\sigma,E,\alpha}f(x) &= \sum_{j_1=0}^{N}\sum_{j_2=0}^{N}\int_{\mathbb{S}^1}\int_{\mathbb{R}^+}[(1-\mathbb{1}_U)|\alpha|^2](z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))w_{j_1}(x,\theta) \\ &\qquad \cdot [w_{j_2}f](z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta),\theta_{j_2-j_1}(x,\theta))\,dt\,d\theta \\ &\quad + \sum_{j_1=0}^{N-1}\sum_{j_2=0}^{N-1}\int_{\mathbb{S}^1}\int_{\mathbb{R}^+}[\mathbb{1}_U|\alpha|^2](z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))w_{j_1}(x,\theta) \\ &\qquad \cdot [w_{j_2}f](z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta),\theta_{j_2-j_1}(x,\theta))\,dt\,d\theta. \end{aligned}\tag{37}$$

Here we've used the fact that $(1-\mathbb{1}_U)^2 = 1-\mathbb{1}_U$. Then break off the part with $j_2 = j_1$ to obtain

$$\begin{aligned} N_{\sigma,E,\alpha,ballistic}f(x) &= \sum_{j=0}^{N}\int_{\mathbb{S}^1}\int_{\mathbb{R}^+}[(1-\mathbb{1}_U)|\alpha|^2](z_{-j}(x,\theta),\theta_{-j}(x,\theta))w_j(x,\theta) \\ &\qquad \cdot [w_jf](x+\tau_-(x,\theta)\theta+t\theta,\theta)\,dt\,d\theta \\ &\quad + \sum_{j=0}^{N-1}\int_{\mathbb{S}^1}\int_{\mathbb{R}^+}[\mathbb{1}_U|\alpha|^2](z_{-j}(x,\theta),\theta_{-j}(x,\theta))w_j(x,\theta) \\ &\qquad \cdot [w_jf](x+\tau_-(x,\theta)\theta+t\theta,\theta)\,dt\,d\theta.. \end{aligned}$$

Now we have to be a bit careful in order to combine both summands. In the first summand, whenever $(z_{-m}(x,\theta),\theta_{-m}(x,\theta))$ is in the support of $(1-\mathbb{1}_U)\alpha^2$, we know that $x+\tau_-(x,\theta)\theta+t\theta$ is a line segment very near the corner c where f vanishes. Thus we can throw out the term involving $j=m$. Now for the terms with $j<m$, we may immediately combine the summands into one, thus eliminating the presence


$(R_{1,1}(x), S_{1,1}^{-1}(\theta))$
(x, θ)
$(q_1(x,\theta), S_{1,1}^{-1}(\theta))$

FIGURE 4. Here we demonstrate the construction of $q_1(x,\theta)$ in the case of a beam of broken rays very close to a corner point.

of $\mathbb{1}_U$. That is, we have

$$\begin{aligned} N_{\sigma,E,\alpha,ballistic}f(x) = &\sum_{j=0}^{m-1}\int_{\mathbb{S}^1}\int_{\mathbb{R}^+}|\alpha|^2(z_{-j}(x,\theta),\theta_{-j}(x,\theta))w_j(x,\theta) \\ &\cdot [w_j f](x+\tau_-(x,\theta)\theta+t\theta,\theta)\,dt\,d\theta \\ &+\sum_{j_1=m+1}^{N}\int_{\mathbb{S}^1}\int_{\mathbb{R}^+}[(1-\mathbb{1}_U)|\alpha|^2](z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))w_{j_1}(x,\theta) \\ &\cdot [w_{j_1} f](x+\tau_-(x,\theta)\theta+t\theta,\theta)\,dt\,d\theta \\ &+\sum_{j_2=m}^{N-1}\int_{\mathbb{S}^1}\int_{\mathbb{R}^+}[\mathbb{1}_U|\alpha|^2](z_{-j_2}(x,\theta),\theta_{-j_2}(x,\theta))w_{j_2}(x,\theta) \\ &\cdot [w_{j_2} f](x+\tau_-(x,\theta)\theta+t\theta,\theta)\,dt\,d\theta. \end{aligned} \tag{38}$$

For the second and third terms of (38), we use the fact that for each j_2 with $m \leq j_2 \leq N-1$, $x+(\tau_-(x,\theta)+t)\theta$ in the third term has the same reflection signature (l_1, l_2) as in the second term, but with $j_1 = j_2+1$. This has to do with the supports of the corresponding cutoff functions. Fix j_1, j_2 such that $m \leq j_2 = j_1 - 1$, and let (l_1, l_2) be the reflection signature of (x,θ) relative to $(z_{-j_1}(x,\theta), \theta_{-j_1}(x,\theta))$ in the second term or $(z_{-j_2}(x,\theta), \theta_{-j_2}(x,\theta))$ in the third. The key idea is that both reflection signatures will be the same. Recall the circumscribed disk $B((\frac{1}{2}, \frac{1}{2}), \frac{\sqrt{2}}{2})$. Similar to the case with no corner point, we consider the sequence of reflection signatures $\{(h_1, v_1), (h_2, v_2), \ldots, (h_{N-1}, v_{N-1})\}$ in the unfolded domain where all of the unfolded broken rays from U touch and over which f has nonzero integral. Thus we've cut out the tiny "bow-tie" region near the corner c, which is why there are only $N-1$ such pairs. Also recall for $0 \leq m \leq N-1$ the functions $q_m(x,\theta)$ as the *final* intersection point of the line $R_{h_m,v_m}(x) + tS^{-1}_{h_m,v_m}(\theta)$ with ∂B for $t \leq 0$.

Clearly, q_m is smooth on the set of (x, θ) for which the integrand doesn't vanish, since ∂B is smooth (see Figure 4). We then have a local regularized version of the weight function w_j given by

$$\begin{aligned} w_{reg,j}(x,\theta) &= \exp\left(-\int_{\mathbb{R}^+} \sigma(x-\tau\theta,\theta)\,d\tau\right) \\ &\quad\cdot \prod_{m=1}^{j} \exp\left(-\int_{\mathbb{R}^+} \sigma(q_m(x,\theta)+\tau S^{-1}_{h_m,v_m}(\theta), S^{-1}_{h_m,v_m}(\theta))\,d\tau\right), \end{aligned} \tag{39}$$

for $0 \le j \le N-1$. By construction, $w_{reg,j}(x,\theta)$ coincides with $w_j(x,\theta)$ or $w_{j-1}(x,\theta)$ on the broken rays we've isolated, with the index depending on whether the broken ray intersects c. This allows us to rewrite the ballistic part of the normal operator as

$$\begin{aligned} &N_{\sigma,E,\alpha,ballistic} f(x) \\ &= \sum_{j=0}^{m-1} \int_{\mathbb{S}^1}\int_{\mathbb{R}^+} |\alpha|^2(z_{-j}(x,\theta),\theta_{-j}(x,\theta))w_j(x,\theta)[w_j f](x+\tau_-(x,\theta)\theta+t\theta,\theta)\,dt\,d\theta \\ &+ \sum_{j=m}^{N-1} \int_{\mathbb{S}^1}\int_{\mathbb{R}^+} |\alpha|^2(q_j(x,\theta),S^{-1}_{h_j,v_j}(\theta))w_{reg,j}(x,\theta)[w_{reg,j} f](x+\tau_-(x,\theta)\theta+t\theta,\theta)\,dt\,d\theta. \end{aligned}$$

From here, we can make the change of variables $s = t+\tau_-(x,\theta)$. Then from Lemma 2 of [FSU08] we have a classical pseudodifferential operator of order -1.

Now we need to deal with $N_{\sigma,E,\alpha,reflect}$. However, the approach is similar. Recall we have

$$\begin{aligned} N_{\sigma,E,\alpha,reflect} f(x) = &\sum_{j_1=0}^{N}\sum_{j_2=0,j_2\neq j_1}^{N} \int_{\mathbb{S}^1}\int_{\mathbb{R}^+} [(1-\mathbb{1}_U)|\alpha|^2](z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))w_{j_1}(x,\theta) \\ &\cdot [w_{j_2} f](z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta),\theta_{j_2-j_1}(x,\theta))\,dt\,d\theta \\ &+ \sum_{j_1=0}^{N-1}\sum_{j_2=0,j_2\neq j_1}^{N-1} \int_{\mathbb{S}^1}\int_{\mathbb{R}^+} [\mathbb{1}_U|\alpha|^2](z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))w_{j_1}(x,\theta) \\ &\cdot [w_{j_2} f](z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta),\theta_{j_2-j_1}(x,\theta))\,dt\,d\theta. \end{aligned} \tag{40}$$

Again suppose that the broken ray γ_{x_0,θ_0} intersects a corner c at the mth reflection, where $1 \le m \le N-1$. We claim that in the first summation above, the terms with $j_2 = m$ vanish. To see this, note that if (x,θ) is such that $z_{-j_1}(x,\theta), \theta_{-j_1}(x,\theta)$ is in the support of $(1-\mathbb{1}_U)|\alpha|^2$, then reflecting $j_2 = m$ times from that point will put us in the small "bow-tie" region near c on which the corresponding line integral vanishes. So we rewrite the first term in (40) as

$$\begin{aligned} &\sum_{j_1=0}^{N}\sum_{j_2=0,j_2\neq j_1,j_2\neq m}^{N} \int_{\mathbb{S}^1}\int_{\mathbb{R}^+} [(1-\mathbb{1}_U)|\alpha|^2](z_{-j_1}(x,\theta),\theta_{-j_1}(x,\theta))w_{j_1}(x,\theta) \\ &\cdot [w_{j_2} f](z_{j_2-j_1}(x,\theta)+t\theta_{j_2-j_1}(x,\theta),\theta_{j_2-j_1}(x,\theta))\,dt\,d\theta \end{aligned} \tag{41}$$

The important thing is that in (41) the reflection signature of $z_{j_2-j_1}$ relative to z_{-j_1} for each pair (j_1, j_2) is the same wherever $\alpha^2(1-\mathbb{1}_U)$ are supported.

As in §4.2, for each pair of indices (j_1, j_2), we consider the reflection signature when going from (x,θ) to $(z_{j_2-j_1}, \theta_{j_2-j_1})$. This is given by $(h_{j_2}-h_{j_1}, v_{j_2}-v_{j_1})$. So as in (32) we define

$$\begin{aligned}
\widetilde{y}_{j_1,j_2} &= R_{(h_{j_2}-h_{j_1},v_{j_2}-v_{j_1})}(y)\\
\widehat{\eta} &= \frac{1}{|\widetilde{y}_{j_1,j_2}-x|}((-1)^{h_{j_2}-h_{j_1}}(\widetilde{y}^1_{j_1,j_2}-x^1),(-1)^{v_{j_2}-v_{j_1}}(\widetilde{y}^2_{j_1,j_2}-x^2))\\
\widehat{\theta} &= \frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|},
\end{aligned}$$

and we can rewrite

$$\begin{aligned}
&N_{\sigma,E,\alpha,reflect}f(x)\\
&=\sum_{j_1=0}^{N-1}\sum_{j_2=0,j_2\neq j_1}^{N-1}\int_{\mathbb{R}^2}\left[(1-\mathbb{1}_U)|\alpha|^2\right]\left(q_{j_1}\left(x,\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right),S^{-1}_{h_{j_1},v_{j_1}}\left(\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right)\right)\\
&\quad\cdot w_{reg,j_1}\left(x,\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right)w_{reg,j_2}(y,\widehat{\eta})\,f(y)\,\frac{dy}{|\widetilde{y}_{j_1,j_2}-x|}\\
&+\sum_{j_1=0}^{N-1}\sum_{j_2=0,j_2\neq j_1}^{N-1}\int_{\mathbb{R}^2}[\mathbb{1}_U|\alpha|^2]\left(q_{j_1}\left(x,\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right),S^{-1}_{h_{j_1},v_{j_1}}\left(\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right)\right)\\
&\quad\cdot w_{reg,j_1}\left(x,\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right)w_{reg,j_2}(y,\widehat{\eta})\,f(y)\,\frac{dy}{|\widetilde{y}_{j_1,j_2}-x|}\\
&=\sum_{j_1=0}^{N-1}\sum_{j_2=0,j_2\neq j_1}^{N-1}\int_{\mathbb{R}^2}\left|\alpha\left(q_{j_1}\left(x,\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right),S^{-1}_{h_{j_1},v_{j_1}}\left(\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right)\right)\right|^2\\
&\quad\cdot w_{reg,j_1}\left(x,\frac{\widetilde{y}_{j_1,j_2}-x}{|\widetilde{y}_{j_1,j_2}-x|}\right)w_{reg,j_2}(y,\widehat{\eta})\,f(y)\,\frac{dy}{|\widetilde{y}_{j_1,j_2}-x|}.
\end{aligned}\tag{42}$$

Finally, we are ready to prove the result about the decomposition of $N_{\sigma,E,\alpha}$.

Proof of Proposition 2. Recall that we have already shown in the discussion, see (22), that

$$N_{\sigma,E,\alpha}=N_{\sigma,E,\alpha,ballistic}+N_{\sigma,E,\alpha,reflect}.$$

where $N_{\sigma,E,\alpha,ballistic}$ is a classical pseudo differential operator of order -1 that is elliptic on $\mathcal{M}$ (more generally on the subset of the cotangent space given by $\mathcal{M}'$). So we have a parametrix Q of $N_{\sigma,E,\alpha,ballistic}$ of order 1 that is elliptic on $\mathcal{M}$.

For the reflected part, we have a sum of terms of the form (32) in the case of a neighborhood of broken rays that don't have any corner points (except possibly as a starting point), or $N_{\sigma,E,\alpha,reflect}$ has the form of (42), both of which have C_0^∞ Schwartz kernels. Applying Q to both sides of (22) yields the result. ☐

Proposition 3. *Under the conditions of Theorem 2, without assuming that $I_{\sigma,E,\alpha}$ is injective,*

(a) *one has the a priori estimate*

$$\|f\|_{L^2(K)}\leq C\|N_{\sigma,E,\alpha}f\|_{H^1(\Omega)}+C_s\|f\|_{H^{-s}(\Omega)},\quad \forall s;$$

(b) $\operatorname{Ker} I_{\sigma,E,\alpha}$ *is finite dimensional and included in $C^\infty(K)$.*

Proof. The proof is essentially the same as that for ([FSU08], Proposition 3). To prove part (a), by Proposition 2 we have that

$$QN_{\sigma,E,\alpha}f = f + QN_{\sigma,E,\alpha,reflect}f + S_1 f =: f + S_2 f.$$

S_2 has C_0^∞ Schwarz kernel and by elliptic regularity we have

$$\|f\|_{L^2(K)} - C_s\|f\|_{H^{-s}(\Omega)} \leq \|QN_{\sigma,E,\alpha}f\|_{L^2(K)} \leq C\|f\|_{H^1(\Omega)}$$

as desired.

For part (b), if $f \in \operatorname{Ker} I_{\sigma,E,\alpha}$, then $(\mathrm{Id} + S_2)f = 0$, and also S_2 is a compact operator on $L^2(K)$ with smooth kernel. By the Fredholm theorem, f lies in a finite dimensional subspace of $C^\infty(\Omega)$. □

5. Reducing the Smoothness Requirement on σ, α

Given a choice of smooth cutoff α and a smooth σ vanishing near $\partial\Omega$ and such that $I_{\sigma,E,\alpha}$ is injective, we would like to be able to perturb α and σ in C^2 slightly and still have $N_{\sigma,E,\alpha}$ be injective. We do this according to the following modified version of ([FSU08], Proposition 4).

Proposition 4. *Assume that σ, α are fixed and belong to C^2. Let (σ', α') be $O(\delta)$ close to (σ, α) in C^2. Then there exists a constant $C > 0$ that depends on an a priori bound on the C^2 norm of (σ, α) such that*

$$\|(N_{\sigma',E,\alpha'} - N_{\sigma,E,\alpha})f\|_{H^1(\Omega)} \leq C\delta\|f\|_{L^2(K)}. \tag{43}$$

Proof. Recall that $N_{\sigma,E,\alpha} = N_{\sigma,E,\alpha,ballistic} + N_{\sigma,E,\alpha,reflect}$, where $N_{\sigma,E,\alpha,ballistic}$ is a weakly singular integral operator with smooth kernel and $N_{\sigma,E,\alpha,reflect}$ is smoothing. By the discussion in the proof of Proposition 4 of [FSU08], we have

$$\|N_{\sigma,E,\alpha,ballistic} - N_{\sigma',E,\alpha',ballistic}\|_{H^1(\Omega)} \leq C\delta\|f\|_{L^2(K)}. \tag{44}$$

Similarly, the Schwartz kernels of $N_{\sigma,E,\alpha,reflect}$ depend smoothly on σ and α (see (32) and (42)), so the same estimate holds for this term. The completes the proof. □

Proof of Theorem 2. First note that if $I_{\sigma,E,\alpha} : L^2(K) \to L^2(\Gamma_-)$ is injective, then $N_{\sigma,E,\alpha} : L^2(K) \to H^1(\Omega)$ is also injective by an integration by parts. Next, since $N_{\sigma,E,\alpha,ballistic}$ is elliptic on $\mathcal{M}$, we may apply a parametrix Q to get the equation $QN_{\sigma,E,\alpha}f = f + S_1 f + QN_{\sigma,E,\alpha,reflect}f$. Note that $S_1 + QN_{\sigma,E,\alpha,reflect}$ is compact. So by elliptic regularity we have the estimate

$$\|f\|_{L^2(K)} \leq C(\|N_{\sigma,E,\alpha}f\|_{H^1(\Omega)} + \|(S_1 + QN_{\sigma,E,\alpha,reflect})f\|_{L^2(\Omega)}).$$

By Lemma 2 of [SU05], the compactness of the second term on the right implies that

$$\|f\|_{L^2(K)} \leq C'\|N_{\sigma,E,\alpha}f\|_{H^1(\Omega)}.$$

The other inequality is obvious.

Finally, to prove part (b), we just use Proposition 4 to get

$$\begin{aligned}\|f\|_{L^2(K)} &\leq C\|N_{\sigma_0,E,\alpha^0}f\| \\ &\leq C\|N_{\sigma_0,E,\alpha^0}f\|_{H^1(\Omega)} - C\|(N_{\sigma_0,E,\alpha^0} - N_{\sigma,E,\alpha})f\|_{H^1(\Omega)} \\ &\leq C\|N_{\sigma_0,E,\alpha^0}f\|_{H^1(\Omega)} - C^2\delta\|f\|_{L^2(K)}.\end{aligned}$$

□

6. Discussion

In this work, we have utilized the machinery of [FSU08] for the X-ray transform with generic weights and applied it to the given broken ray transform on the Euclidean square. The technique of path unfolding can be applied to any polyhedral domain, however the square is very nice in that reflections are well-defined even at corner points. The unfolding approach also placed significant restraints on the attenuation coefficient σ, which is why we could only reasonably tackle the case $\sigma = 0$ and then use the normal operator computation to extend the results to C_0^2 perturbations. It is perhaps of interest to carry out similar computations for the broken ray transform on higher dimensional polyhedral domains, although it is not clear how one would be able to deal with corners other than to cut them out of the measurements altogether. One interesting problem to consider would be to try to get analogous microlocal results for the unit ball. However, one potential obstacle for strictly convex smooth domains is that a beam of collimated rays, upon reflection becomes uncollimated. This makes it difficult, or perhaps impossible, to use smooth cutoff functions of the measurements to isolate individual beams of broken rays from which one can obtain a nice integral form of the normal operator. Regardless, it is the author's hope that more progress can be made on this particular problem and its variants in the future.

References

[Bal04] G. Bal. On the attenuated radon transform with full and partial measurements. *Inverse Problems*, 20:399–419, 2004.

[BS04] J. Boman and J.O. Strömberg. Novikov's inversion formula for the attenuated radon transform - a new approach. *The Journal of Geometric Analysis*, 14(2), 2004.

[CS96] M. Choulli and P. Stefanov. Inverse scattering and inverse boundary value problems for the linear Boltzmann equation. *Comm. Partial Differential Equation*, 21(5-6):763–785, 1996.

[CS98] M. Choulli and P. Stefanov. An inverse boundary value problem for the stationary transport equation. *Osaka J. Math.*, 36(1):87–104, 1998.

[Esk04] G. Eskin. Inverse boundary value problems in domains with several obstacles. *Inverse Problems*, 20(5):1497–1516, 2004.

[Fin86] D.V. Finch. Uniqueness for the attenuated x-ray transform in the physical range. *Inverse Problems*, 2:197–203, 1986.

[Fin04] D.V. Finch. *The attenuated X-ray transform: Recent developments*. MSRI Publications. Cambridge University Press, Cambridge, UK, 2004. in Inside Out, Inverse Problems, and Applications, G. Uhlmann, ed.

[FSU08] B. Frigyik, P. Stefanov, and G. Uhlmann. The x-ray transform for a generic family of curves and weights. *J. Geom. Anal.*, 18(1):81–97, 2008.

[GS94] A. Grigis and J. Sjöstrand. *Microlocal Analysis for Differential Operators: An Introduction*, volume 196 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, New York, 1994.

[H8̈3] L. Hörmander. *The Analysis of Linear Partial Differential Operators*, volume 1-3. Springer-Verlag, New York, 1983.

[Hub11] M. Hubenthal. An inverse source problem in radiative transfer with partial data. *Inverse Problems*, 27(12), 2011.

[Ilm12] J. Ilmavirta. Broken ray tomography in the disk. arXiv:1210.4354 [math-ph], 2012.

[KS12] C. Kenig and M. Salo. The calderón problem with partial data on manifolds and applications. arXiv:1211.1054 [math.AP], 2012.

[Nat01] F. Natterer. Inversion of the attenuated radon transform. *Inverse Problems*, 17:113–119, 2001.

[Nov02a] R.G. Novikov. An inversion formula for the attenuated x-ray transformation. *Ark. Mat.*, 40(1):145–167, 2002.

[Nov02b] R.G. Novikov. On the range characterization for the two-dimensional attenuated x-ray transformation. *Inverse Problems*, 18, 2002.

[Qui93] E.T. Quinto. Singularities of the x-ray transform and limited data tomography in $\mathbb{R}^2$ and $\mathbb{R}^3$. *SIAM J. Math. Anal.*, 24:1215–1225, 1993.

[Qui05] E.T. Quinto. An introduction to x-ray tomography and radon transforms. In *Symposia in Applied Mathematics*, 2005.

[Ste08] P. Stefanov. Microlocal approach to tensor tomography and boundary and lens rigidity. *Serdica Math. J.*, 34(1):67–112, 2008. An extended version of the mini-course I gave at the Symposium on Inverse Problems, Rio De Janiero, Jan. 2007.

[SU04] P. Stefanov and G. Uhlmann. Stability estimates for the x-ray transform of tensor fields and boundary rigidity. *Duke Math. J.*, 123:445–467, 2004.

[SU05] P. Stefanov and G. Uhlmann. Boundary rigidity and stability for generic simple metrics. *J. Amer. Math. Soc.*, 18(4):975–1003, 2005.

[SU08] P. Stefanov and G. Uhlmann. An inverse source problem in optical molecular imaging. *Analysis and PDE*, 1(1):115–126, 2008.

[SU12] P. Stefanov and G. Uhlmann. The geodesic x-ray transform with fold caustics. *Analysis and PDE*, 5(2):219–260, 2012.

[Tab] S. Tabachnikov. Billiards. http://www.math.psu.edu/tabachni/Books/billiardsbook.pdf.

[Tab05] S. Tabachnikov. *Geometry and Billiards.* American Mathematical Society, 2005.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ
E-mail address: `john.m.hubenthal@jyu.fi`